

1 Homogeneous Linear ODE's with Constant Coefficients

Homogeneous, linear ODEs with constant coefficients were likely covered in your Differential Equations course (MTH 256 or equiv.). If you need a review, please see:

Constant Coefficients, Homogeneous

or your differential equations text.

Answer the following questions for each differential equation below:

- identify the order of the equation,
- find the number of linearly independent solutions,
- find an appropriate set of linearly independent solutions, and
- find the general solution.

Each equation has different notations so that you can become familiar with some common notations.

(a) $\ddot{x} - \dot{x} - 6x = 0$

(b) $y''' - 3y'' + 3y' - y = 0$

(c) $\frac{d^2w}{dz^2} - 4\frac{dw}{dz} + 5w = 0$

2 Inhomogeneous Linear ODEs with Constant Coefficients (First Example)

Inhomogeneous, linear ODEs with constant coefficients are among the most straightforward to solve, although the algebra can get messy. This content should have been covered in your Differential Equations course (MTH 256 or equiv.). If you need a review, please see: The Method for Inhomogeneous Equations or your differential equations text.

The general solution of the **homogeneous** differential equation

$$\ddot{x} - \dot{x} - 6x = 0$$

is

$$x(t) = A e^{3t} + B e^{-2t}$$

where A and B are arbitrary constants that would be determined by the initial conditions of the problem.

(a) Find a particular solution of the inhomogeneous differential equation $\ddot{x} - \dot{x} - 6x = -25 \sin(4t)$.

(b) Find the general solution of $\ddot{x} - \dot{x} - 6x = -25 \sin(4t)$.

(c) Some terms in your general solution have an undetermined coefficients, while some coefficients are fully determined. Explain what is different about these two cases.

(d) Find a particular solution of $\ddot{x} - \dot{x} - 6x = 12e^{-3t}$

(e) Find the general solution of $\ddot{x} - \dot{x} - 6x = 12e^{-3t} - 25 \sin(4t)$

How is this general solution related to the particular solutions you found in the previous parts of this question?

Can you add these particular solutions together with arbitrary coefficients to get a new particular solution?

(f) Sense-making: **Check your answer;** Explicitly plug in your final answer in part (e) and check that it satisfies the differential equation.

3 Shifted Sinusoids

None A fun and fascinating fact about sinusoids is that a (normalized) sum of a cosine and a sine function is just a shifted cosine function. You can see this relationship geometrically using the Geogebra applet in <https://books.physics.oregonstate.edu/GMM/fouriergg.html>. Below is an algebraic proof:

For a and b related by the normalization condition $a^2 + b^2 = 1$, we have

$$a \cos \theta + b \sin \theta = a \frac{e^{i\theta} + e^{-i\theta}}{2} + b \frac{e^{i\theta} - e^{-i\theta}}{2i} \quad (1)$$

$$= \frac{a - ib}{2} e^{i\theta} + \frac{a + ib}{2} e^{-i\theta} \quad (2)$$

$$= \frac{1}{2} e^{-i\phi} e^{i\theta} + \frac{1}{2} e^{i\phi} e^{-i\theta} \quad (3)$$

$$= \frac{1}{2} e^{i(\theta-\phi)} + \frac{1}{2} e^{-i(\theta-\phi)} \quad (4)$$

$$= \cos(\theta - \phi) \quad (5)$$

where in (1) we have used Euler's formula, in (2) we have regrouped terms, in (3) we have rewritten $a + ib = e^{i\phi}$, in (4) we have combined exponents, and in (5) we have used the inverse form of Euler's formulas.

In the rest of this class, we will discuss 4 different forms of the solutions of the equation of motion for a simple harmonic oscillator:

A Form: $x(t) = A \cos(\omega t + \phi)$

B Form: $x(t) = B_1 \cos(\omega t) + B_2 \sin(\omega t)$

C Form: $x(t) = C e^{i\omega t} + C^* e^{-i\omega t}$

D Form: $x(t) = \operatorname{Re}[D e^{i\omega t}]$

- (a) In the calculation above, identify lines of the calculation that represent the A, B, and C forms.
- (b) From your identifications, find relationships between the constants in the A, B, and C forms.